

EVOLUTION OF THE FUCACEAE (PHAEOPHYCEAE) INFERRED FROM nrDNA-ITS<sup>1</sup>

*Ester A. Serrão,<sup>2</sup> Lawrence A. Alice,<sup>3</sup> and Susan H. Brawley<sup>4</sup>*

Department of Plant Biology and Pathology, University of Maine, 5722 Deering Hall, Orono, Maine 04469-5722

Sequences of the internal transcribed spacer region (ITS-1, 5.8S, and ITS-2) of nuclear ribosomal DNA were obtained from 16 species representing all six genera of Fucaceae (*Ascophyllum*, *Fucus*, *Hesperophycus*, *Pelvetia*, *Pelvetiopsis*, and *Xiphophora*) plus one outgroup (*Hormosira*). Parsimony analysis indicated that the family Fucaceae is monophyletic and that the northern hemisphere taxa are highly divergent from the only southern hemisphere genus, *Xiphophora*. The genus *Pelvetia* is not monophyletic because the European *P. canaliculata* is more closely related to *Fucus*, *Hesperophycus*, and *Pelvetiopsis* than to other *Pelvetia* species. We establish *Silvetia*, gen. nov. and transfer the 3 Pacific species of *Pelvetia* to the new genus. *Fucus* is monophyletic and not ancestral in the Fucaceae. The ITS sequences identified two strongly supported lineages within *Fucus*, one with *F. serratus* sister to the clade containing *F. gardneri*, *F. distichus*, and *F. evanescens* and a second including *F. vesiculosus*, *F. spiralis*, *F. ceranoides*, and *F. virsoides*. The ITS was not useful for resolving relationships within each of these clusters and between populations of *F. vesiculosus*. Within-individual




FIG. 1. Speciation of the northern hemisphere Fucaceae, as proposed by Powell (1963) and represented here in cladogram format. This hypothesis was based on the characters 1) monoecy/dioecy and 2) the number of eggs produced per oogonium (this is indicated between brackets at the branches where each state originated). Following the initial divergence of the dioecious and monoecious clades, we represented a trichotomy in each clade because Powell did not discuss the divergence hierarchy at this level.

relationships in Fucaceae has been widely followed, it is phylogenetically invalid because the proposed divergence of Fucaceae into monoecious and dioecious lineages would prevent *Fucus* from being monophyletic. This leads to two mutually exclusive questions: 1) is *Fucus* monophyletic, or 2) are the Fucaceae divided into two lineages corresponding to monoecy and dioecy? The evolutionary relationships within the genus *Pelvetia* are also controversial, as earlier suggestions (Serrão 1996, Serrão and Brawley 1997) and recent work by Lee et al. (1998) have suggested that *P. canaliculata* might form a separate lineage distinct from the Pacific *Pelvetia*.

The nuclear rDNA internal transcribed spacers (ITS-1 and ITS-2) are used widely in phylogenetic studies at the species level (reviews by Hamby and Zimmer 1992, Baldwin et al. 1995). They have been used extensively in algae (e.g. Kooistra et al. 1992, Coleman et al. 1994, Goff et al. 1994, Bakker et al. 1995a, b, van Oppen et al. 1995), brown algae in particular (Saunders and Druehl 1993, van Oppen

et al. 1993, Peters et al. 1997, Stache-Craigne et al. 1997), and most recently in *Fucus* (Leclerc et al. 1998).

In this study, we use ITS sequences to investigate phylogenetic relationships among the six genera of the Fucaceae as well as species relationships within *Fucus* and *Pelvetia*. We also examine Powell's hypothesis for the evolution of monoecy/dioecy and number of eggs per oogonium in the Fucaceae in light of our ITS-based phylogeny.

#### MATERIALS AND METHODS

**Sample material.** Most species of *Fucus* occur in the Atlantic except *F. gardneri*, a Pacific endemic, and *F. virsoides*, an endemic of the Mediterranean Sea. In addition to their Atlantic distributions, *F. spiralis* occurs in the eastern Pacific and *F. evanescens* in the western Pacific. All species of *Pelvetia* occur in the Pacific except *Pelvetia canaliculata*. *Ascophyllum* is monotypic and occurs exclusively in the Atlantic. *Pelvetiopsis* and *Hesperophycus* are also monotypic, and they are found exclusively in the Pacific. *Xiphophora* occurs in the Australasian region and is the only genus from the southern hemisphere in the Fucaceae. All currently recognized species of each Fucaceae genus were sampled (Table 1) with the exception of *Xiphophora gladiata*, which occurs only in Tasmania. The ITS sequences of each species were determined from several locations when possible. Voucher specimens from most of these locations were deposited in the University of Maine's herbarium.

When it was possible to obtain reproductive material, sperm were used for the DNA extractions because they yielded DNA in greater quantities and better quality (i.e. they have lower polysaccharide contamination and are free of endophytes, such as the fungus that occurs in *A. nodosum* and *P. canaliculata*). To obtain uncontaminated sperm cells, receptacles were washed with distilled water and placed in petri dishes under the light. As they dried, antheridia (containing sperm) were expelled through the pore of each conceptacle in an orange, mucilaginous mass. A few of these were picked from just outside the pore of each conceptacle with a sterile needle at a dissecting microscope, frozen in liquid N<sub>2</sub>, and stored at -280°C until extraction. When it was not possible to obtain fresh sperm cells, vegetative tips from fresh, herbarium, or silica-gel-dried material were used. Material from the Baltic Sea was ground in the CTAB extraction buffer (see following discussion) and stored at room temperature for approximately 1 month before extraction.

**DNA extraction.** DNA was extracted using a CTAB method (modified after Doyle and Doyle 1987). Antheridia were ground in microcentrifuge tubes with 2X CTAB buffer (2% CTAB, 1.4 M NaCl, 0.2% β-mercaptoethanol, 20 mM EDTA, 100 mM Tris HCl pH 8, 1% PVP-40), and vegetative tips were ground in liquid N<sub>2</sub> before adding the CTAB buffer. The samples were incubated with 2X CTAB buffer and proteinase K at 60°C for 5 h with frequent agitation, followed by two purification steps with chloroform:isoamyl alcohol and precipitation with isopropanol. RNA in the redissolved pellet was digested and RNase removed with phenol plus chloroform:isoamyl alcohol followed by an overnight precipitation with 100% ethanol and ammonium acetate. DNA quality and quantity in the redissolved pellet were assessed on agarose gels and microfluorometrically.

**PCR amplification.** The ITS region of each sample was symmetrically amplified by PCR either with the primers ITS5 and ITS4 (White et al. 1990; Fig. 2) or with F5 and F4 (Fig. 2), which we specifically designed for *Fucus* from a sequence for the SSU of rDNA available for *F. gardneri* at GenBank (X53987) and on our own sequences for the 5' end of the LSU of *Fucus*. Amplification conditions were 1 X buffer II, 1.9 mM MgCl<sub>2</sub>, 0.2 mM/dNTP, 0.3 mM/primer, 1.25 U of Stoffel Fragment or AmpliTaq (Perkin Elmer, Norwalk, Connecticut), and 2 ng of genomic DNA per 25-μL reaction. The PCR parameters were 94°C for 5 min, 45 cycles of 97°C for 1 min, 48°C for 2 min, and 72°C for 3 min, followed by a final extension step at 72°C for 15 min. These conditions

TABLE 1. List of sampled species of the Fucaceae used to sequence the ITS region. In addition to these, the five ITS sequences from Leclerc et al. (1998) were also used in the global analyses. GenBank accession numbers in boldface represent the sequences selected for the more extended analyses. Multiple numbers for the same geographic region represent ITS sequences determined from a single individual, except in the case of *Hesperophycus californicus*, for which the first two sequences are from one individual and the other two sequences are from another individual.

Species	Geographic origin	GenBank accession number	University of Maine Herbarium accession number
<i>Ascophyllum nodosum</i> (Linnaeus) Le Jolis	Isle of Man (Port St. Mary Ledges)	AF102971	—
	Maine coast (Schoodic Point, Acadia National Park)	AF102961, AF102962, AF102963	UMAL278 <sup>a</sup>
	Maine estuary (Penobscot River, near Bucksport)	<b>AF102964</b>	UMAL279 <sup>a</sup>
	Massachusetts (Woods Hole)	AF102965, AF102966	UMAL271
	Northern Portugal (Viana do Castelo, Minho)	AF102969	UMAL274
	Norway (Sotra, near Bergen)	AF102967	UMAL258
	Western Ireland (Galway)	AF102968	—
	White Sea (Ivanov Navolok, Russia)	AF102970	UMAL256
	Isle of Man (Castletown Harbour)	AF102903	UMAL266
	<i>Fucus ceranoides</i> Linnaeus	Maine coast (Chamberlain)	AF102935
White Sea (Ivanov Navolok, Russia)		AF102936	UMAL272
<i>Fucus distichus</i> Linnaeus	Japan (Muroran, Hokkaido)	AF102937	—
	Maine coast (Schoodic Point, Acadia National Park)	AF102938	UMAL276 <sup>a</sup>
<i>Fucus evanescens</i> C. Agardh	White Sea (Ivanov Navolok, Russia)	AF102939	UMAL252
	California (Pigeon Point, San Mateo)	<b>AF102940</b>	UMAL265
<i>Fucus gardneri</i> P.C. Silva	Washington (San Juan Island)	AF102941	UMAL259
	Nova Scotia (Lunenburg Cove)	<b>AF102943</b> , AF102944, AF102945	—
<i>Fucus serratus</i> Linnaeus	Norway (Sotra, near Bergen)	AF102942	UMAL260
	Canary Islands (Punta de Galdar, Gran Canaria)	AF102904	UMAL268
	Maine coast (Schoodic Point, Acadia National Park)	AF102905-6	UMAL280 <sup>a</sup>
	Northern Portugal (Viana do Castelo, Minho)	AF102907	UMAL262
	Oregon (Boiler Bay)	AF102908	—
<i>Fucus vesiculosus</i> Linnaeus	Washington (San Juan Island)	AF102909	UMAL269
	Algarve, Portugal (Ria Formosa Natural Park, near Faro)	AF102910	UMAL257
	Central Baltic Sea (Askö, near Stockholm)	AF102911, AF102912, <b>AF102913</b>	—
	Isle of Man (Port St. Mary)	AF102919-21	UMAL261
	Maine estuary (Penobscot river, near Bucksport)	AF102926-7	UMAL281 <sup>a</sup>
	Maine coast (Schoodic Point, Acadia National Park)	AF102922	UMAL282
	Northern Baltic Sea (Drivan, near Umeå)	AF102914-8	UMAL255
	Northern Portugal (Viana do Castelo, Minho)	AF102925	UMAL283
	Skagerrak (Tjärnö, Swedish west coast)	AF102928-9	—
	Southern Norway (Sotra, near Bergen)	AF102923-4	UMAL253
<i>Fucus virsoides</i> J. Agardh	White Sea (Ivanov Navolok, Russia)	AF102930-2	UMAL273
	Mediterranean (Trieste, Adriatic Sea)	AF102933	—
<i>Hesperophycus californicus</i> P.C. Silva	Mediterranean (Venice, Adriatic Sea)	AF102934	—
	California (between Lovers' Point and Point Piños, Monterey)	AF102946, <b>AF102947</b> , AF102948, AF102949, AF102950	UMAL251, UMAL250
<i>Pelvetia babingtonii</i> (Harvey) De Toni	Japan (Muroran, Hokkaido)	<b>AF102957</b>	UMAL254
<i>Pelvetia canaliculata</i> (Linnaeus) Decaisne et Thuret	Isle of Man (Port St. Mary, Chapel Bay)	AF102955	UMAL264
	Northern Portugal (Viana do Castelo, Minho)	AF102954	UMAL263
<i>Pelvetia compressa</i> (J. Agardh) De Toni	Southern Norway (Sotra, near Bergen)	<b>AF102953</b>	UMAL267
	California (Pigeon Point, San Mateo)	<b>AF102956</b>	UMAL275 <sup>a</sup>
<i>Pelvetia siliquosa</i> Tseng et C.F. Chang	China (Shandong Peninsula)	<b>AF102958</b> , AF102959, AF102960	—
<i>Pelvetiopsis limitata</i> (Setchell) Gardner	California (Pigeon Point, San Mateo)	<b>AF102951</b>	UMAL270
	Oregon (Boiler Bay)	AF102952	—
<i>Xiphophora chondrophylla</i> (R. Brown ex Turner) Montagne ex Harvey	Southern Australia (Flinders)	<b>AF102972</b>	—
<i>Hormosira banksii</i> (Turner) Decaisne (5 outgroup)	Southern Australia (Sorrento)	<b>AF102973</b> , AF102974, AF102975, AF102976, <b>AF102977</b> , AF102978, AF102979	—

<sup>a</sup> A morphologically similar specimen from the same site.




Fig. 2. The nrDNA ITS region showing the relative position of the 39 end of the SSU gene, ITS-1, the 5.8S gene, ITS-2, the 59 end of the LSU gene, and the location of the primers used. Sequences for primers F1 through F5 are shown; primers ITS3 and ITS4 are from White et al. (1990). Primers not to scale. \* 5 primers used in PCR amplification.

were optimized to amplify preferentially the ITS region from the Fucaceae over that from endophytes, which was always smaller.

PCR products were electrophoresed in agarose gels and run long enough to isolate the target 1000–1100-bp ITS band of the Fucaceae from other, always smaller fragments that were present in some cases, such as the ITS region of *Mycosphaerella ascophylli* Le Jolis, an endophytic fungus in *Ascophyllum nodosum* and *Pelvetia canaliculata*. Sequencing of these smaller bands followed by GenBank searches confirmed that they belonged to the genus *Mycosphaerella* (data not shown), whereas GenBank searches matching the sequences of the 1000–1100-bp products confirmed that these belonged to the Phaeophyceae. The target bands were cut under UV light, and the agarose was digested with  $\lambda$ -agarase by the manufacturer's instructions (FMC, Rockville, ME). These products were either cloned or used for the sequencing reactions without further purification.

**Cloning.** To determine the extent of intraindividual polymorphism in some taxa and/or populations, the whole ITS region was cloned (TOPO TA cloning kit, Invitrogen Corp., La Jolla, California) following the manufacturer's instructions, and plasmid DNA from each recombinant colony was extracted using alkaline lysis minipreps with PEG precipitation (Applied Biosystems Inc., User Bulletin #18).

**Sequencing.** The ITS region of one to five individuals of each species or population was sequenced in both directions using the Sanger dideoxy chain termination method for cycle sequencing with dye-labeled terminators (Perkin Elmer) on an ABI 373 automated sequencer (Applied Biosystems, Foster City, California). For direct sequencing, the primers used to sequence the ITS-2 region were ITS3 or F3 and ITS4 or F4 (Fig. 2). The primers used to sequence the ITS-1 region were ITS5 or F5, F2, or in a few cases F1. Where the whole ITS region had been cloned, it was possible to fully sequence from the 39 end of the SSU gene to the 59 end of the LSU gene in both directions (over 1000 bp) using universal primers in the vector.

**Alignment.** The chromatogram output for each sample was edited using the software Sequence Navigator 1.0.1 (Applied Biosystems Inc.), and the sequences were manually aligned. To determine the ITS boundaries for the Fucaceae, the flanking sequences of ITS-1 and ITS-2 were aligned with all ITS sequences listed in GenBank that include the flanking sequence for the SSU, the 5.8S, and the LSU genes. These sequences represent a wide range of organisms, including other protists, animals, fungi, and plants. The limits of ITS-1 and ITS-2 varied among all these sequences; thus, the positions at which a sharp change occurs from a sequence that is conserved across all taxa to a sequence that becomes variable (and vice versa for the 39 ends of the ITS regions) are those most likely to be the real boundaries between the rRNA genes and the spacers, and these were considered as such in this study. Of the 82 sequences used, 77 were determined in this study

from several geographic regions and in some cases from several clones within individuals. This alignment is available at GenBank. An additional five sequences from Leclerc et al. (1998) from Roscoff, Brittany, were included in this study, but we considered *F. lutarius* to be a form of *F. vesiculosus* (as in Powell 1963).

**Outgroup selection.** The sister group to the Fucaceae is unknown because different morphological characteristics suggest different evolutionary patterns. Our choice was based on the suggestion by Clayton (1984) that the families of Fucales radiated from Australasia because all but one family of Fucales occur in Australasia and all are represented by species with primitive characteristics. In the Fucaceae, *Xiphophora* is the Australasian genus with primitive characteristics. *Hormosira* is an Australasian taxon that shares unique characteristics with *Xiphophora* (see Clayton 1984) and was thus used as outgroup. In addition to this, we also obtained full sequences (data not shown) for other Fucales: *Cystoseira*, *Bifurcaria*, and *Cystosphaera* (the latter was once included in the Fucaceae), but these were almost unalignable.

**Phylogenetic analyses.** Given the large number of sequences and poor resolution within species, it was impossible within reasonable computational time to execute a heuristic search to completion with all sequences. For this reason, a bootstrap (1000 replicates) consensus tree was determined for 82 Fucaceae ITS sequences using the "fast" stepwise addition method, as implemented in PAUP 4.0b1 (Swofford 1998), to select representative sequences for a more comprehensive analysis. The basal-most sequence was selected from each species or each clade of poorly resolved species because this sequence is the one most likely to have the plesiomorphic states as opposed to derived apomorphic sequences in terminal clades. Character state changes were all weighted equally. Polymorphisms were treated as uncertain, and insertions/deletions (indels) were coded as missing data to avoid including them as if they were many independent events while still retaining the information about substitutions in other taxa in the indel regions. In a separate analysis, indels were assumed to be single evolutionary events and were given the same weight as nucleotide substitutions by coding them in a separate matrix as binary characters. Large indels and those from regions where these tend to occur frequently by strand slippage (e.g. microsatellite regions) were not included in the analysis because they were unlikely to be the result of single events and positional homology could not be accurately determined.

Phylogenetic inferences, using the selected subset of sequences, used three different optimality criteria: maximum parsimony, maximum likelihood, and neighbor joining. Maximum-likelihood analysis was based on the Hasegawa-Kishino-Yano model. Neighbor-joining analyses were performed with three distance models: uncorrected distances, Jukes-Cantor, and Kimura 2-parameter. Inferences based on Fitch parsimony used the Branch-and-Bound algorithm, which guarantees finding the shortest trees (Hillis et al. 1993). All analyses were performed with PAUP 4.0b1. To test for phylogenetic signal in the data, we determined the index of skewness ( $g_1$ ) of the distribution of tree lengths of 100,000 random trees using PAUP and compared it to the critical value for 15 taxa and 500 variable characters in Hillis and Huelsenbeck (1992). Confidence limits of individual clades were estimated by bootstrap analysis (Felsenstein 1985) with 100,000 replicates, and decay analysis (Bremer 1988, Donoghue et al. 1992) was performed with AutoDecay 3.0 (Eriksson and Wikstrom 1996) and the reverse constraint option in PAUP. Two primary characters used in Powell's classification were mapped onto the ITS tree using MacClade 3.0 (Maddison and Maddison 1992).

## RESULTS

**ITS properties and alignment.** In the Fucaceae, the length of ITS-1 varied between 401 and 515 bp and that of ITS-2 between 322 and 409 bp (Table 3). We were unable to obtain the ITS-2 sequence from *X. chondrophylla*. These length values are not comparable to those published for *Fucus* and other brown algae (Saunders and Druehl 1993, van Oppen et al.

TABLE 2. Length and GC content of ITS-1 and ITS-2 region in the Fucaceae and *Hormosira banksii* (n.d. 5 not determined).

Taxon	ITS-1	
	Length (bp)	GC (%)

TABLE 4. Ranges of pairwise distances (%) in the ITS region between 1) species of each genus, 2) geographic regions for each species, 3) individuals in each region, and 4) ITS fragments cloned from single individuals. Sequences from Leclerc et al. (1998) were included in this analysis, but we considered *Fucus lutarius* to be a form of *Fucus vesiculosus* (as in Powell 1963). *Xiphophora* is not included in the table because only one sequence was determined. (n 5 number of species, regions/species, individuals/region or clones/individual involved, respectively, where more than one were determined; n.a. 5 not applicable; n.d. 5 not determined).

Genus	Between species	Between regions (geographic)	Within regions	Within individuals
<i>Ascophyllum</i>	n.a.	0.0–2.2 (n 5 9)	n.d.	0.0–0.2 (n 5 2–3)
<i>Fucus</i>	0.1–7.2 (n 5 8)	0.0–5.6 (n 5 2–11)	0.3 (n 5 2)	0.1–2.6 (n 5 2–5)
<i>Hesperophycus</i>	n.a.	n.d.	0.2–0.3 (n 5 2)	0.0–0.5 (n 5 2–3)
<i>Pelvetia</i>	0.8–13.3 (n 5 4)	0.0 (n 5 2)	n.d.	0.0 (n 5 3)
<i>Pelvetiopsis</i>	n.a.	0.2 (n 5 2)	n.d.	n.d.
<i>Hormosira</i>	n.a.	n.d.	n.d.	0.0–0.8 (n 5 7)

well-supported clades of *Fucus*, but the only clearly monophyletic species in this genus is *F. serratus*. The ITS sequences are not useful to resolve most relationships between the species within each of these two clusters. *Fucus serratus* is a sister group to a clade containing *F. gardneri*, *F. distichus*, and *F. evanescens*, none of which appears to be monophyletic, and the relationships within the clade containing *F. vesiculosus*, *F. spiralis*, *F. virsoides*, and *F. ceranoides* are not resolved. Thus, the basal-most sequence for *F. serratus* was selected for further analyses, and the remaining *Fucus* species were represented by a sequence from only one species from each of the other two clades. The same tree topology was obtained if each of the selected sequences was replaced by any other sequence from its clade or by the two most divergent ones in the clade.

The phylogenetic analyses based on the reduced data set yielded one most parsimonious tree of length 768 (Fig. 4), and this was consistent with the tree obtained with all sequences (Fig. 3). The distribution of tree lengths is significantly skewed to the left ( $g1 \ 5 \ 21.26$ ), indicating that the data have phylogenetic signal (Hillis 1991, Hillis and Huelsenbeck 1992). The data show low homoplasy, as indicated by a high consistency index (0.90) and retention index (0.89). The ITS tree reveals several clades, most of which are well supported by high bootstrap and decay values (Fig. 4). The data support the Fucaceae as a monophyletic group. In particular, the inclusion of the only southern hemisphere genus of this family, *Xiphophora*, in the same clade as the other Fucaceae is well supported (Fig. 4). The results were identical when the analysis was performed with most indels coded as binary (presence/absence) characters. Maximum-likelihood and neighbor-joining analyses also resulted in the same tree topology as with parsimony, the only exception being the clade with the three Pacific *Pelvetia*, which collapses into an unresolved trichotomy when either the Kimura 2-parameter or the Jukes-Cantor model is used.

Within the Fucaceae of the northern hemisphere, *A. nodosum* is sister to all other genera (Fig. 4). *Pelvetia* is not monophyletic because its Atlantic species, *P. canaliculata*, shares a common ancestor with

the genera *Hesperophycus*, *Pelvetiopsis*, and *Fucus*, thus being more closely related to these than to the other *Pelvetia*. Imposing a constraint tree to force monophyly of *Pelvetia* adds 13 steps to the most parsimonious tree. However, the Pacific species of *Pelvetia* (*P. babingtonii*, *P. compressa*, and *P. siliquosa*) are monophyletic, as is *Fucus*. All clades are well supported except the node that separates *A. nodosum* from all other northern Fucaceae. This was an unresolved node when all sequences were included in the bootstrap analysis (Fig. 3); the alternative possibility would be uniting *A. nodosum* with the Pacific *Pelvetia* in a monophyletic clade.

#### DISCUSSION

The Fucaceae appear monophyletic, although *Xiphophora*, the only southern hemisphere genus, is more distantly related to other Fucaceae than all northern Fucaceae are among themselves. Recent studies using rDNA SSU and LSU sequences to study the Fucaceae did not find enough variability to resolve their phylogenetic relationships (Rousseau et al. 1997, Lee et al. 1998). *Ascophyllum nodosum* appears basal for the northern hemisphere Fucaceae, contradicting suggestions that it might be more related to *Fucus* than to *Pelvetia canaliculata*, although these were poorly supported (Rousseau et al. 1997). On the basis of ITS sequences, the Pacific *Pelvetia* are the closest group to *A. nodosum*.

*Pelvetia* does not appear monophyletic. Thus, *P. canaliculata* needs to be recognized as a monotypic genus, creating a monophyletic clade including only the Pacific species of *Pelvetia* under a new genus (see below), as *P. canaliculata* is the type species for *Pelvetia*. Additional molecular data support this change (Lee et al. 1998). It is interesting to note that both of the most primitive Fucaceae of the Atlantic Ocean, *A. nodosum* and *P. canaliculata*, share the characteristic of being the hosts of an endophytic species of fungus (*Mycosphaerella ascophylli*) that is absent in all other Fucaceae (Kohlmeyer and Kohlmeyer 1979).

The idea of an ancestral *Fucus* that gave rise to all other Fucaceae (Powell 1963) is not consistent with our results or with those of Leclerc et al. (1998), who used ITS sequences to compare five species of *Fucus* from Brittany and suggested that these were

63	—	<i>F. ceranoides</i> - Isle of Man
	—	<i>F. ceranoides</i> - Roscoff
	—	<i>F. lutarius</i> - Roscoff
	—	<i>F. spiralis</i> - Roscoff
56	—	<i>F. vesiculosus</i> - Roscoff
	—	<i>F. spiralis</i> - Maine coast
	—	<i>F. spiralis</i> - Maine coast
84	—	<i>F. spiralis</i> - Oregon
	—	<i>F. spiralis</i> - Oregon




FIG. 3. Bootstrap consensus tree, including all ITS sequences for the Fucaceae. Taxa in bold were selected to represent each species or clade in further analyses. Sequences with the same label represent multiple clones from the same individual except in the case of *Hesperophycus californicus*, for which the first two sequences are from one individual and the other three from another one. *Hormosira banksii* is the outgroup.

the result of a very recent radiation because sequence variability was very low. However, our results are in agreement with Leclerc et al. (1998) in suggesting a very recent radiation for *Fucus*, although for opposite reasons: a recent rapid radiation is a possible explanation for the high within-individual variability that we observed in *Fucus*. Also, our rooted ITS tree for the Fucaceae shows *Fucus* as the most derived genus. These results are also consistent with Leclerc et al. (1998) in that *F. serratus* is basal to the remaining *Fucus* species, although none of the spe-

cies that our study indicates as sister to *F. serratus* was sampled in their study.

*Fucus* is monophyletic, and all species of *Fucus* are closely related. The well-supported (Fig. 3) clustering of *F. spiralis* (monoecious), *F. virsoides* (monoecious), *F. vesiculosus* (dioecious), and *F. ceranoides* (dioecious) indicates that these are closely related and contradicts expectations that all monoecious versus dioecious species of *Fucus* diverged early in the evolutionary history of the Fucaceae (Powell 1963; Fig. 1). Further evidence against an early di-




FIG. 4. The single most parsimonious tree for the ITS region of the taxa selected for the final analyses. Bootstrap (above) and decay (below) support values for the clades are indicated.

vergence of dioecious versus monoecious species comes from the fact that *F. serratus*, which is dioecious and has thus been assumed to be more closely related to *F. vesiculosus*, appears in the same lineage that gave rise to monoecious species: *F. distichus*, *F. evanescens*, and *F. gardneri*. Our results (Fig. 3) suggest that these might not be monophyletic species. These were once classified as a single species with several subspecies that differed in morphology and habitat (Powell 1957). Although *F. distichus* and *F. evanescens* are very closely related, they are reproductively isolated to a large degree, at least in New England, because their reproductive seasons do not overlap to a great extent and especially because reproduction in *F. distichus* occurs in tide pools at low tide (Pearson and Brawley 1996), when the chance of cross-fertilization with the low-shore species *F. evanescens* is reduced. *Fucus gardneri* might be reproductively isolated from these because it occurs only in the eastern Pacific Ocean, although ITS sequences suggest that *F. evanescens* from Japan might be more related to *F. gardneri* than to Atlantic *F. evanescens*.

The character "monoecious/dioecious" has




FIG. 5. Evolution of the characters (A) monoecy/dioecy and (B) number of eggs per oogonium in the Fucaceae as inferred from sequences of the ITS region.

changed several times during the evolutionary history of the Fucaceae (Fig. 5A), and the same was found in Desmarestiaceae (Peters et al. 1997). This suggests that in brown algae, monoecy/dioecy can be changed by relatively simple mechanisms and thus is not a good indicator of evolutionary relationships. The monoecious condition might have been more favorable to rapid, long-distance migration of small populations because the Fucaceae do not produce vegetative propagules. For effective dispersal of dioecious species that reproduce only sexually, it is necessary that enough individuals of both sexes are present for sexual reproduction to occur with the high rates of success that have been observed in fu-



coid populations (Brawley 1992, Pearson and Brawley 1996, Serrão et al. 1996).

The phylogeny based on ITS sequences does not support the hypothesis (Fritsch 1945, Powell 1963; Fig. 1) that producing eight eggs per oogonium is the most primitive state in Fucaceae. However, this character seems to have evolved mostly in the direction of a progressive reduction in the number of viable eggs produced per oogonium, *Fucus* being the only exception (Fig. 5B). The inclusion of *Hesperophycus* and *Pelvetiopsis* in the same clade suggests that the character state "producing one egg per oogonium" evolved only once, in the common ancestor to these genera, also contradicting Powell's (1963) hypothesis (Fig. 1). Thus, although the character egg number does not follow the trend suggested by Powell (1963), it seems to be a useful character to distinguish the clades revealed by the ITS data.

The placement of *Xiphophora* as sister to the northern hemisphere Fucaceae is supported by ITS data. *Xiphophora* has unusual features for the Fucaceae, and Clayton (1984) has suggested that its evolutionary history should be further investigated. *Xiphophora* lacks clearly defined receptacles, a characteristic in common with *Hormosira*. The sperm of *Xiphophora* have different features from the sperm of northern hemisphere Fucaceae (Clayton 1994). Instead, they share structural similarities with the monotypic families Hormosiraceae and Himanthaliaceae (Clayton 1994), especially the possession of a spine on the anterior flagellum of the sperm (Moestrup 1982). This has been used as a basis to propose that the genera *Xiphophora*, *Hormosira*, and *Himanthalia* might be related (Moestrup 1982), a hypothesis that warrants further investigation of the evolutionary history of *Himanthalia*. On the other hand, the fucacean characteristics of *Xiphophora* are dichotomous branching and a four-sided apical cell (Clayton 1984). Sperm ultrastructure (Manton 1964, 1965) suggests that the Cystoseiraceae are a primitive state and that *Fucus* represents the most advanced condition. *Hormosira* (Hormosiraceae) seems related to *Xiphophora* (Fucaceae) on the basis of sperm ultrastructure (Manton 1965) and the absence of true receptacles but not on the basis of other characteristics. However, the character "apical cell" suggests that the Seirococcaceae might be the most related family to the Fucaceae, as both have a four-sided apical cell (Clayton 1984), although another character (one single mature egg produced per oogonium) suggests that the Seirococcaceae might be more closely related to the Cystoseiraceae instead (Clayton 1984). Given this controversy, we attempted to include representatives of all three of these families (*Hormosira* (Hormosiraceae), *Cystosphaera* (Seirococcaceae), and *Cystoseira* and *Bifurcaria*, both Cystoseiraceae) in the outgroup, but only *Hormosira* was alignable in the ITS region, suggesting that it is the closest relative of Fucaceae.

The intraindividual variability in the ITS region of *Fucus* species and the occurrence of shared inter-specific polymorphisms in each of the two unresolved clusters of *Fucus* species might be a consequence of gene flow between species through frequent hybridization and/or rapid recent radiation at a rate that exceeds the homogenization rate by concerted evolution (Hillis and Davis 1988, Williams et al. 1988). The latter is consistent with the fact that low intraindividual variability was found in *A. nodosum*, a basal genus. The hybridization hypothesis is supported by the fact that sequences from an individual *F. vesiculosus* might be more similar to ITS sequences from *F. spiralis* than to other sequences from that same individual. *Fucus* species can be cross-fertilized in the laboratory (Thuret 1854, Williams 1899, Kniep 1925, Burrows and Lodge 1953, Bolwell et al. 1977, Edwards et al. 1997), and numerous reports exist of the occurrence of putative *Fucus* hybrids in nature (Sauvageau 1909, Stomps, 1911, Kniep 1925, Burrows and Lodge 1951, Scott and Hardy 1994). The only species in this cluster that cannot hybridize with the others in nature is the Mediterranean-endemic *Fucus virsoides* because it is geographically isolated from all other Fucaceae.

How can these results help explain the bipolar distribution of the Fucaceae? Also, if the Fucaceae originated in southern Australia, where the Fucales seem to have evolved and diversified (Clayton 1984), when and through which route did they colonize the northern Pacific and Atlantic Oceans while remaining absent from South America and most of Africa (except for *Fucus* in northern Morocco and the Canary Islands)?

If sequence divergence and divergence time are correlated in this data set, the large divergence in ITS sequences (Table 2) between the Fucaceae of the southern and northern hemispheres indicates that these have been isolated for a long period. An ancient separation of these cold-adapted seaweeds could be the result of 1) a transequatorial crossing during a colder period or 2) cold adaptation and consequent hemispheric divergence of an ancestor that might have been initially adapted to the warmer conditions of lower latitudes. The first of these hypotheses seems to be more likely because it is consistent with the hypothesis that the order Fucales evolved and radiated during the Mesozoic on the southern shores of Australasia (Clayton 1984). This is supported by the fact that genera from all but one (Himanthaliaceae) of the families of Fucales occur there, and those have characteristics that are considered primitive in each family (Clayton 1984). Thus, the most likely origin for the evolution of the Fucaceae might be in Australia, where a Fucaceae ancestor common to *Xiphophora* and to the northern Fucaceae might have been able to colonize or drift across the equatorial fringe, invading the Pacific, during a cooler period. A similar southern hemisphere ancient origin followed by a transequatorial

crossing is suggested by ITS sequences in the Desmarestiaceae (Peters et al. 1997).

Clayton (1984) suggests that the opportunity for the early fucoids to migrate to the northern hemisphere might have been provided by the Pleistocene glaciations (2 Myr to 18,000 years ago), when the ocean's temperatures had decreased significantly and a land bridge existed between Australia and Indonesia (Clayton 1984). However, the large sequence distances between *Xiphophora* and the northern Fucaceae suggest that their divergence would have started before this period, perhaps during the Oligocene/Miocene (38–7 Myr ago). In comparison, sequence divergence was very low in the ITS region of Arctic and Antarctic species of a brown algal genus hypothesized to have been isolated since the Pleistocene glaciations (van Oppen et al. 1993). Alternatively, the levels of divergence observed in this DNA region of the Fucaceae might not be proportional to the evolutionary time since divergence. This could have been the case if rapid speciation events occurred in a short time, when possibly small populations of Fucaceae might have spread through the very hostile and thus very selective intertidal regions.

After the northern Fucaceae became isolated from the southern hemisphere ones, ancestors to *A. nodosum* and *P. canaliculata* might have invaded the Atlantic, where they acquired the endophytic fungus *Mycosphaerella*. However, if this took place through crossing the Arctic, it would have been possible only 3–3.5 Myr ago, when the Bering Strait opened (e.g. Lüning 1990, Vermeij 1991, Dunton 1992)<sup>5</sup>. The opening of the Bering Strait, which had been closed for approximately 60 Myr, was followed by several events of opening and closing of this connection between the Pacific and the Atlantic by way of the Arctic. These events might have been involved in the origin of the divergence between ancestors to *Fucus* and *Hesperophycus/Pelvetiopsis* and later between Pacific and Atlantic *Fucus*. The western Atlantic shores lack *P. canaliculata* and *F. serratus* is present only in a restricted area, probably as a species introduced there from Europe by humans in the 1800s (Robinson 1903, Edelstein et al. 1971–73). These shores might have been secondarily colonized by the Fucaceae following the glaciated periods when cold-adapted rocky shore species, forced to migrate south, might have found suitable rocky intertidal shores on the eastern Atlantic but not on the western Atlantic (Lüning 1990, Palumbi and Kessing 1991). Secondary recolonization of the western Atlantic following the Pleistocene glaciations has also

been suggested for other macroalgae (e.g. Lüning 1990, van Oppen et al. 1995).

The geographic distribution of all but one of the *Fucus* species in the Atlantic and its adjacent seas, and the fact that the only Pacific-endemic species (*F. gardneri*) is not basal for the genus, suggests that *Fucus* evolved in the northern Atlantic Ocean and only later colonized the Pacific. Thus, the evolution of *Fucus gardneri* in the Pacific might be associated with a transarctic passage of an Atlantic ancestor common to *F. serratus* and *F. gardneri*, although most transarctic crossings documented for marine animals (e.g. Palumbi and Kessing 1991, Vermeij 1991) and macroalgae (e.g. Stam et al. 1988, van Oppen et al. 1994, 1995) occurred from the Pacific to the Atlantic. *Fucus distichus* and *F. evanescens*

<sup>5</sup> Note in proof: L. Marinovich and A. Y. Gladenkov (1999, Evidence for an early opening of the Bering Strait. *Nature* 397: 149–151) present data in support of the first opening of the Bering Strait occurring between 4.8–7.4 Myr. This is in agreement with our sequence data suggesting that the colonization of the Atlantic by the Fucaceae took place earlier than the data previously estimated for the first opening of the Bering Strait.

the results suggest that some of these might not be monophyletic and/or that hybridization might occur between some species within each cluster. 7) During evolution of the Fucaeeae, the character number of eggs per oogonium has not followed a gradual reduction. The character monoecy/dioecy has changed several times, thus having little value in defining evolutionary relationships in the family.

We transfer *Pelvetia babingtonii* (Harvey) De Toni, *P. compressa* (J. Agardh) De Toni, and *P. siliquosa* Tseng et C. F. Chang to *Silvetia*, gen. nov. This transfer is based upon anatomical differences and our molecular analysis of the species presently included in *Pelvetia*.

**Silvetia** E. Serrão T. O. Cho, S. M. Boo et Brawley, gen. nov.

Diagnosis: Novum genus Fucaeeae plerumque ovis cum duobus per oogonium. Verbum ipsum Pelvetiae simile est, sed etiam dissimile putandum est propter divisionem quandam oogoniorum potius in longitudinem vel in obliquitatem quam ex transverso perductam; propter varietatem quandam in ipso ordine quo aciditas nuclei cum signis de rDNA inter se transcriptis complectitur.

New genus of the Fucaeeae, typically with 2 eggs per oogonium, similar in morphology to *Pelvetia* but distinguished from it by cleavage of the oogonia being longitudinal or oblique, as opposed to transverse; by differences in the nucleic acid sequence of the internal transcribed spacers (ITS-1, ITS-2) of the rDNA.

Type species: **Silvetia compressa** (J. Agardh) E. Serrão, T. O. Cho, S. M. Boo et Brawley, comb. nov.

Basionym: *Furoidium compressum* J. Agardh (*Sp. Alg.* 1:204. 1848)

Etymology: Named in honor of Paul C. Silva, a keen student of the furoid algae and Curator of Algae at the Herbarium of the University of California at Berkeley.

We also propose the following new combinations:

**Silvetia babingtonii** (Harvey) E. Serrão, T. O. Cho, S. M. Boo et Brawley, comb. nov.

Basionym: *Fucus babingtonii* Harvey (*Proc. Amer. Acad. Arts & Sci.* 4:329. 1860).

**Silvetia siliquosa** (Tseng & C. F. Chang) E. Serrão, T. O. Cho, S. M. Boo et Brawley, comb. nov.

Basionym: *Pelvetia siliquosa* Tseng & C. F. Chang (*Acta Bot. Sin.* 2:293. 1953).

The oogonia of the species that we transfer to *Silvetia* typically cleave longitudinally or slightly obliquely (Yendo 1907, Gardner 1910, Tseng and Chang 1953), rather than transversely as in *Pelvetia canaliculata* (e.g. Fritsch 1945). Cleavage is irregular if more than the typical 2 eggs per oogonium are produced (Gardner 1910, Tseng and Chang 1983,

Song et al. 1996). Similar abnormalities in egg number and cleavage are observed, in a small number of oogonia in most Fucaeeae (pers. obs.). These variations may result from defective cleavage and/or an abnormal targeting of the number of haploid nuclei retained in the young oogonium (see Fritsch 1945). Both *Pelvetia* and *Silvetia* include perennial fucaean algae that lack a midrib and are found in the upper intertidal zone, but those taxa transferred here to *Silvetia* tend to be more cartilaginous than *Pelvetia*. Ecotypic factors contribute to variations in how cartilaginous the various species of *Silvetia* are and to their stature (Yoshida and Silva 1992, Song et al. 1996). Silva (1996) includes a photograph (his Fig. 3, showing *Fucus compressus* C. Agardh) of an isotype of *Furoidium compressum* and provides related discussion. Molecular phylogenetic analysis has demonstrated that the oogonial plane of division is a robust character (cf. Yendo 1907).

Advice from Dr. Debashish Bhattacharya was invaluable to us. We are also grateful for helpful comments on an earlier draft by Dr. Irv Kornfield, to Dr. Jeanine Olsen, and for the comments by two anonymous reviewers and Dr. Wytze Stam that greatly improved the manuscript. Many thanks to Dr. Christopher Campbell for use of his equipment, to Patty Singer and Dr. Diane Caporale for running the automated sequencer, to Dr. Gareth Pearson for general help, and to all who collected and mailed specimens: Dr. Kathryn van Alstyne (Oregon), Dr. Christine Ashburner (Australia), Dr. Jon Ashen (California), Dr. Mauro Bastianini (Venice), Dr. Carolyn Bird (Nova Scotia), Dr. Vanda Brotas (South Portugal), Dr. Ian Davison (Antarctic), Dr. Megan Dethier (Washington State), Dr. D. L. Duan (China), Ben Hale (California), Dr. Ricardo Haroun (Canary Islands), Jenny Hedman (Norway and northern Sweden), Dr. Joanna Jones (Isle of Man), Dr. Lena Kautsky (southern Sweden), Dr. Erin Krellwitz (North Carolina), Rhoda Lin (California), Marina Lipizer (Trieste), Dr. Arthur Mathieson (New Hampshire), Joe McIntyre (northern Sweden), Dr. Taizo Motomura (Japan), Dr. Isabel Pinto (northern Portugal), and Dr. Rui Santos (Algarve) and Dr. Richard Searles (North Carolina), Dr. Dagmar Stengel (Galway), Brian Wysor (Long Island), and the staff of the Kartesh Marine Station of the University of St. Petersburg for help during collecting trips in the White Sea, Russia. We appreciate nomenclatural advice from Drs. Michael Guiry, Christine Maggs, Richard Moe and an anonymous referee and thank Estelle Sheehan for the Latin diagnosis. We appreciate the collaboration of Tae Oh Cho and Dr. Sung Min Boo (Chungnam National University) in the diagnosis; they will present additional work elsewhere. This study was supported by the NSF grant OCE-9216981 to S.H.B. and by a bequest of H. Franklin Brawley. E.A.S. is also thankful for graduate fellowships from the Fulbright Commission, JNICT (Portugal), the Luso-American Foundation, and the Center for Marine Studies-Sea Grant of Maine/New Hampshire (U.S.A.).

Bakker, F. T., Olsen, J. L. & Stam, W. T. 1995a. Evolution of nuclear rDNA ITS sequences in the *Cladophora albida/sericea* Clade (Chlorophyta). *J. Mol. Evol.* 40:640-51.

Bakker, F. T., Stam, W. T. & Olsen, J. L. 1995b. Global phylogeography in the cosmopolitan species *Cladophora vagabunda* (Chlorophyta) based on nuclear rDNA ITS sequences. *Eur. J. Phycol.* 30:197-208.

Baldwin, B. G., Sanderson, M. J., Porter J. M., Wojciechowski, M. F., Campbell, C. S. & Donoghue, M. J. 1995. The ITS region of nuclear ribosomal DNA: a valuable source of evidence on angiosperm phylogeny. *Ann. Missouri Bot. Garden* 82:247-77.

- Bolwell, G. P., Callow, J. A., Callow, M. E. & Evans, L. V. 1977. Cross-fertilization in fucoid seaweeds. *Nature* 268:626-7.
- Brawley, S. H. 1992. Fertilization in natural populations of the dioecious brown alga *Fucus ceranoides* L. and the importance of the polyspermy block. *Mar. Biol.* 113:145-57.
- Bremer, K. 1988. The limits of amino acid sequence data in angiosperm phylogenetic reconstruction. *Evolution* 42:795-803.
- Burrows, E. M. & Lodge, S. M. 1951. Autecology and the species problem in *Fucus*. *J. Mar. Biol. Assoc. UK* 30:161-76.
- 1953. Culture of *Fucus* hybrids. *Nature* 172:1009-10.
- Clayton, M. N. 1984. Evolution of the Phaeophyta with particular reference to the Fucales. In Round, F. E. & Chapman, D. J. [Eds.] *Progress in Phycological Research*, vol. 3. Biopress, Bristol, pp. 11-46.
- 1994. Circumscription and phylogenetic relationships of the southern hemisphere family Seirococcaceae (Phaeophyceae). *Bot. Mar.* 37:213-20.
- Coleman, A. W., Suarez, A. & Goff, L. J. 1994. Molecular delimitation of species and syngens in Volvocacean green algae (Chlorophyta). *J. Phycol.* 30:80-90.
- Decaisne, J. & Thuret, G. 1845. Recherches sur les antheridies et les spores de quelques *Fucus*. *Ann. Sci., Ser. 3, Bot.* 3:1-15.

- based on DNA sequences of rRNA-ITS. *Phycologia* 36(Suppl.): 101.
- Serrão, E. A., Pearson, G. A., Kautsky, L. & Brawley, S. H. 1996. Successful external fertilization in turbulent environments. *Proc. Natl. Acad. Sci. USA* 93:5286–90.
- Silva, P. C. 1996. California seaweeds collected by the Malaspina Expedition, especially *Pelvetia* (Fucales, Phaeophyceae). *Madroño* 43:345–54.
- Song, H. S., Seo, K. S. & Boo, S. M. 1996. Field studies of the brown alga *Pelvetia siliquosa* with implications for taxonomy and distribution. *Algae* 11:65–71.
- Stache-Crain, B., Müller, D. G. & Goff, L. J. 1997. Molecular systematics of *Ectocarpus* and *Kuckuckia* (Ectocarpales, Phaeophyceae) inferred from phylogenetic analysis of nuclear- and plastid-encoded DNA sequences. *J. Phycol.* 33:152–68.
- Stam, W. T., Bot, P. V. M., Boele-Bos, S. A., van Rooij, J. M. & van den Hoek, C. 1988. Single-copy DNA-DNA hybridizations among five species of *Laminaria* (Phaeophyceae): phylogenetics and biogeographic implications. *Helgoländer Meeresunters.* 42:251–67.
- Stomps, T. J. 1911. Études topographiques sur la variabilité des *Fucus vesiculosus* L., *platycarpus* Thur., et *ceranoides* L. *Recueil l'Institut Bot 'Leo Errera' Brux.* 8:326–77.
- Swofford, D. L. 1998. PAUP (Phylogenetic Analysis Using Parsimony), Version 4.0b1. Sinauer, Sunderland, Massachusetts.
- Thuret, G. 1854. Recherches sur la fécondation des Fucales. *Ann. Sci. Nat. Bot. série 4*, 2:197–214.
- Tseng, C. K. & C. F. Chang. 1953. On a new species of *Pelvetia* and its distribution. *Acta Bot. Sinica* 2:280–97.
- van Oppen, M. J. H., Olsen, J. L., Stam, W. T., Hoek, C. van den & Wiencke, C. 1993. Arctic-Antarctic disjunctions in the benthic seaweeds *Acrosiphonia arcta* (Chlorophyta) and *Desmarestia viridis/willii* (Phaeophyta) are of recent origin. *Mar. Biol.* 115:381–6.
- van Oppen, M. J. H., Diekmann, O. E., Wiencke, C., Stam, W. T. & Olsen, J. L. 1994. Tracking dispersal routes: phylogeography of the Arctic-Antarctic disjunct seaweed *Acrosiphonia arcta* (Chlorophyta). *J. Phycol.* 30:67–80.
- van Oppen, M. J. H., Draisma, S. G. A., Olsen, J. L. & Stam, W. T. 1995. Multiple trans-Arctic passages in the red alga *Phycodrys rubens*: evidence from nuclear rDNA ITS sequences. *Mar. Biol.* 123:179–88.
- Vermeij, G. J. 1991. Anatomy of an invasion: the trans-Arctic interchange. *Paleobiology* 17:281–307.
- White, T. J., Burns, T., Lee, S. & Taylor, J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In Innis, M., Gelfand, D., Sninsky, J. & White, T. [Eds.] *PCR Protocols: A Guide to Methods and Applications*. Academic Press, San Diego, pp. 315–22.
- Williams, J. L. 1899. New *Fucus* hybrids. *Ann. Bot.* 13:187–8.
- Williams, S. C., DeBry, R. W. & Feder, J. L. 1988. A commentary on the use of ribosomal DNA in systematic studies. *Syst. Zool.* 37:60–2.
- Wynne, M. J. & Magne, F. 1991. Concerning the name *Fucus muscoides* (Cotton) J Feldmann et Magne. *Cryptogamie, Algol.* 12: 55–65.
- Yendo, K. 1907. The Fucales of Japan. *J. Coll. Sci. Imp. Univ. Tokyo* 21:1–174.
- Yoshida, T. & P. C. Silva. 1992. On the identity of *Fucus babingtonii* Harvey (Fucales, Phaeophyta). *Japan. J. Phycol.* 40:121–24.